

NOTE

Enumeration of Labelled (k, m) -Trees

Oleg Pikhurko*

*Department of Pure Mathematics and Mathematical Statistics, Cambridge University,
Cambridge CB2 1SB, United Kingdom*E-mail: op201@cam.ac.uk*Communicated by the Managing Editors*

Received February 12, 1998

A k -graph is called a (k, m) -tree if it can be obtained from a single edge by consecutively adding edges so that every new edge contains $k - m$ new vertices while its remaining m vertices are covered by an already existing edge. We prove that there are

$$\frac{(e(k-m)+m)!(e\binom{k}{m}-e+1)^{e-2}}{e!m!((k-m)!)^e}$$

distinct vertex labelled (k, m) -trees with e edges. © 1999 Academic Press

The notion of a tree and its different extensions to k -graphs, that is, k -uniform set systems, play an important role in discrete mathematics and computer science. We will dwell upon the following, rather general, definition suggested independently by Dewdney [5] and Beineke and Pippert [2].

Let us agree that the vertex set is $[n] = \{1, \dots, n\}$. Fix the *edge size* k and the *overlap size* m , $0 \leq m \leq k-1$. We refer to k -subsets and m -subsets of $[n]$ as *edges* and *laps* respectively. A non-empty k -graph without isolated vertices is called a (k, m) -tree if we can order its edges, say E_1, \dots, E_e , so that for every i , $2 \leq i \leq e$, there is i' , $1 \leq i' < i$, such that $|E_i \cap E_{i'}| = m$ and $(E_i \setminus E_{i'}) \cap (\bigcup_{j=1}^{i-1} E_j) = \emptyset$. In other words, we start with a single edge and can consecutively affix a new edge along an m -subset of an existing edge.

Thus, a (k, m) -tree with e edges has $n = e(k-m) + m$ vertices and its edges cover $f = e(\binom{k}{m} - 1) + 1$ laps. For example, a $(k, 0)$ -tree consists of disjoint edges.

* Supported by an External Research Studentship, Trinity College, Cambridge, UK.

The problem of counting $(m+1, m)$ -trees which are known in the literature as m -trees, received great attention and was completely settled by Beineke and Pippert [1] and Moon [9]. This extends celebrated Cayley's theorem [3] as, clearly, 1-trees correspond to usual (Cayley) trees. Later, different bijective proofs for m -trees appeared as well, see [11, 7, 8, 6, 4].

In this paper we enumerate (k, m) -trees. It is not surprising that many of the above counting techniques are applicable here as is, for example, Foata's bijection [7] (details are available from the author [10]). Also, as observed by the referee, our formula can be obtained via Chen's method [4]. We decided to present here an inductive proof which is the shortest one, perhaps.

Let $T_{km}(e)$ denote the number of (k, m) -trees on $[n]$ with e edges, $n = e(k - m) + m$, and let $R_{km}(e)$ count the trees rooted at the lap $[m]$, that is, those trees for which $[m]$ is covered by some edge.

THEOREM 1. *Given integers k, m, e with $0 \leq m \leq k - 1$ and $e \geq 1$, let $n = e(k - m) + m$, $l = \binom{k}{m}$ and $f = e(l - 1) + 1$. Then the number of different (k, m) -trees on $[n]$ equals*

$$T_{km}(e) = \frac{n! f^{e-2}}{e! m! ((k - m)!)^e}. \quad (1)$$

Proof. Like in Beineke and Pippert [1], to prove the theorem, we write down a recurrence relation for $T_{km}(e)$ and then verify that (1) does satisfy the relation. Let us agree that $T_{km}(0) = R_{km}(0) = 1$.

Counting in two different ways the number of pairs (H, L) , where H is a (k, m) -tree on $[n]$ rooted at an m -subset L of $[n]$, we obtain

$$\binom{n}{m} R_{km}(e) = f \cdot T_{km}(e). \quad (2)$$

Next, consider the following method for constructing trees. Select an edge E , a k -subset of $[n]$, and label by L_1, \dots, L_l the laps of E . Represent $e - 1$ as a sum of l non-negative integers, $e - 1 = e_1 + \dots + e_l$. Partition $[n] \setminus E$ into sets X_1, \dots, X_l of sizes $e_1(k - m), \dots, e_l(k - m)$ respectively. On each $L_i \cup X_i$ build a (k, m) -tree H_i rooted at L_i , $i \in [l]$. Clearly, the union of all H_i 's plus the edge E forms a (k, m) -tree with e edges and every such tree H is obtained exactly e times. Therefore, by (2) we obtain

$$\begin{aligned} eT_{km}(e) &= \binom{n}{k} \sum_{\mathbf{e}} \frac{(n - k)!}{(e_1(k - m))! \cdots (e_l(k - m))!} \prod_{i=1}^l R_{km}(e_i) \\ &= \frac{n!}{k!} \sum_{\mathbf{e}} \prod_{i=1}^l \frac{m!(e_i(l - 1) + 1) T_{km}(e_i)}{(e_i(k - m) + m)!}, \end{aligned} \quad (3)$$

where $\sum_{\mathbf{e}}$ denotes the summation over all representations $e-1 = e_1 + \cdots + e_l$ with non-negative integers summands.

Clearly, formula (1) gives correct values for $e=0$. Also, the substitution of (1) into the both sides of (3) gives (after routine cancellations)

$$l(e(l-1)+1)^{e-2} = \sum_{\mathbf{e}} \frac{(e-1)!}{e_1! \cdots e_l!} \prod_{i=1}^l (e_i(l-1)+1)^{e_i-1}. \quad (4)$$

The last identity (in slightly different notation) was established by Beineke and Pippert [1, Lemma 2], which proves our theorem by induction. ■

COROLLARY 1. *The number of labelled m -trees on n vertices, $n > m \geq 1$, is*

$$T_{m+1, m}(n-m) = \binom{n}{m} (mn - m^2 + 1)^{n-m-2}.$$

ACKNOWLEDGMENTS

The author is grateful to the anonymous referee for helpful comments

REFERENCES

1. L. W. Beineke and R. E. Pippert, The number of labelled k -dimensional trees, *J. Combin. Theory* **6** (1969), 200–205.
2. L. W. Beineke and R. E. Pippert, On the structure of (m, n) -trees, in “Proc. 8th Southeast Conference on Combinatorics, Graph Theory and Computing (Louisiana State Univ., Baton Rouge),” *Congressus Numerantium*, Vol. XIX, pp. 75–80, Utilitas Math., Winnipeg, 1977.
3. A. Cayley, A theorem on trees, *Quart. J. Math.* **23** (1889), 376–378.
4. W. Y. C. Chen, A coding algorithm for Rényi trees, *J. Combin. Theory Ser. A* **63** (1993), 11–25.
5. A. K. Dewdney, Multidimensional tree-like structures, *J. Combin. Theory B* **17** (1974), 160–169.
6. O. Egecioğlu and L.-P. Shen, A bijective proof for the number of labelled q -trees, *Ars Combin.* **25** (1988), 3–30.
7. D. Foata, Enumerating k -trees, *Discrete Math.* **1** (1971), 181–186.
8. C. Greene and G. A. Iba, Cayley’s formula for multidimensional trees, *Discrete Math.* **13** (1975), 1–11.
9. J. W. Moon, The number of labeled k -trees, *J. Combin. Theory* **6** (1969), 196–199.
10. O. Pikhurko, Foata’s bijection for (k, m) -trees, manuscript.
11. C. Rényi and A. Rényi, The Prüfer code for k -trees, in “Combinatorial Theory and Its Applications” (P. Erdős et al., eds.) pp. 945–971, North-Holland, Amsterdam, 1970.